

**WEST**

Generate Collection

L15: Entry 7 of 10

File: USPT

Aug 4, 1987

DOCUMENT-IDENTIFIER: US 4684615 A

TITLE: Stabilized isoenzyme control products

## BSPR:

Isoenzymes, or isozymes, as they are alternatively referred to, are enzymes in multiple forms which are capable of performing the same general function but at different rates. They are sufficiently different in chemical composition so that they are generally separable electrophoretically. One such isoenzyme, lactate dehydrogenase (LDH) is found in five electrophoretically distinct fractions. Each of these electrophoretic species of LDH is a tetramer consisting of two polypeptide chain units, H and M, present in different proportion: H.sub.4, MH.sub.3, M.sub.2 H.sub.2, HM.sub.3, and M.sub.4. These five isoenzymes differ in catalytic activity (affinity for the substrate, pyruvate as measured by the Michaelis constant), amino acid composition, heat lability, and immunological responses. The two peptides H and M are coded by different genes. Thus the type of enzyme present is under genetic control and regulated by the conditions of the environment imposed upon the cell.

## BSPR:

In accordance with the objects and principles of the present invention, isoenzyme control reagents are provided for creatinine kinase, lactate dehydrogenase, alanine aminotransferase and aspartase aminotransferase which are substantially stabilized by the addition of plexiform stabilizing means. The preferred plexiform stabilizing means is selected from the group consisting of maltose, mannitol, cellobiose and lactose with the latter most being the most preferred. The plexiform stabilizing means is advantageously provided in a final concentration range of about 2%-8% with the ideal concentration occurring at about 6%. The ideal isoenzyme control reagent will have substantially all water removed, such as by lyophilization, to assist in long term storage and stability.

DEPR:

Enzymes which can be stabilized as described here include oxidoreductases, transferases, dehydrogenases, transaminases, peptidases, hydrolaser, isomerases, mutases and the like. In addition to enzymes already described, other enzymes which could be stabilized herein include but are not limited to malate dehydrogenase; creatine kinase (CK, CPK); alkaline phosphatase (AP, ALP, ALK-phos); aspartate aminotransferase (AST, OT, GOT); alanine aminotransferase (ALT, PT, SGPT); gamma glutamyl transpeptidase (gammaGT, gammaGTP); glutamic oxalacetic transaminase (SGOT); glutamic pyruvic transaminase (GTP); alpha amylase; beta amylase; lactate dehydrogenase (LD, LDH, lactic dehydrogenase); glucose-6-phosphate dehydrogenase (G6PDH); hexokinase (HK); glucose dehydrogenase; glucose oxidase; peroxidase (HRP, HPO, HRPO, PO); glycerol dehydrogenase; glutamate dehydrogenase, cholesterol oxidase; cholesterol esterase; lipase; uricase; urease; glycerol kinase; aldolases; synthetases; nucleases; polymerases; and the like.

DEPR:

The polymer herein can also be used to synthesize peptide bonds. Typically, equimolar solutions containing modified amino acid(s) are dissolved in a solvent system containing an organic solvent such as dioxane or THF (tetrahydrofuran), or a tertiary amine such as (iPr).sub.2 NEt, to form an amino acid mixture. More preferably, the organic solvent is acetonitrile. The solutions dissolved in the solvent system are defined herein as amino acid acceptor and amino acid donor solutions. Examples of such solutions can be found in Table 2 of Example 10.

DEPR:

A conjugated enzyme such as protease (the enzyme incorporated into one of the compounds of the invention) may be added to the amino acid mixture in an amount of about 10.sup.-6 to about 10.sup.-1 mole fraction relative to the amount of amino acid. The conjugated enzyme can be prepared as in Example 8. The reaction is performed at from about 0.degree. to 70.degree. C., at about 1 to about 24 hours. The enzyme polymer conjugate is removed from the formed peptide by filtering and the solvent is evaporated to obtain the resulting peptide.

DETL:

TABLE 2

Enzyme (Solvent, Time) Acceptor Donor Isolated (HPLC) at 23.degree. C. Amino Acid Amino Acid Product Yield (%)

CPC-CT AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH.sub.2 97(100) (Dioxane, 12 h) CPC-CT  
 \* AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH2 94(98) (THF, 12 h) CPC-BPN'  
 CbzLeuLeu-OMe Leu-O.sup.t Bu CbzLeuLeuLeu-O.sup.t Bu 68(95) (Acetonitrile, 22  
 h) CPC-BPN' CbzLeuLeu-OMe PheLeu-O.sup.t Bu CbzLeuLeuPheLeu-O.sup.t Bu 92(98)  
 (Acetronitrile, 24 h) CPC-BPN' CbzValLeu-OMe PheLeu-O.sup.t Bu  
 CbzValLeuPheLeu-O.sup.t Bu 95(98) (Acetronitrile, 24 h) CPC-BPN' CbzValLeu-OMe  
 Ala-NH.sub.2 CbzValLeuAla-NH.sub.2 65(95) (Acetronitrile, 24 h) CPC-BPN'  
 CbzLeuLeu-OMe Ala-NH.sub.2 CbzLeuLeuAla-NH.sub.2 55(90) (Acetronitrile, 24 h)  
 CPC-T CbzPhe-OH Leu-OMe CbzPheLeu-OMe 65(95) (Acetronitrile, 24 h)

at 50.degree. C. \*

**WEST**

Generate Collection

Print

L8: Entry 67 of 71

File: USPT

May 22, 1984

DOCUMENT-IDENTIFIER: US 4450232 A

TITLE: Incorporation of pyridoxal phosphate in dry analytical elements for the determination of enzymes

Brief Summary Text (2):

The present invention relates to dry analytical elements which are useful in determining the amount of pyridoxal phosphate activated enzyme in a liquid sample. The elements of the present invention are particularly useful in quantitating aspartate aminotransferase and alanine aminotransferase enzymes.

Brief Summary Text (4):

Transaminase enzymes are enzymes which catalyze the transfer of an .alpha.-amino group from an .alpha.-amino acid to an .alpha.-keto acid. These enzymes are also sometimes referred to as aminotransferases. Two of the most clinically significant aminotransferase enzymes are L-alanine:.alpha.-ketoglutarate aminotransferase, EC 2.6.1.2 (commonly referred to as alanine aminotransferase or ALT) and L-aspartate:.alpha.-ketoglutarate aminotransferase, EC 2.6.1.1 (commonly referred to as aspartate aminotransferase or AST) (including mitochondrial and cytoplasmic isoenzymes).

Brief Summary Text (11):

Because of the importance of pyridoxal phosphate on the activation of the apo-enzyme form of AST and ALT, this activation has been widely studied. For example, it is known that in dilute solution, activation of the apo-enzyme occurs at low concentrations of pyridoxal phosphate. Thus, the IFCC recommends that the incubation be carried out in a 0.1 mmole per liter pyridoxal phosphate solution. Other studies have indicated that somewhat higher levels of pyridoxal phosphate might be desirable, for example, up to 0.3 mmole per liter. However, in solution, additional pyridoxal phosphate gave little, if any, further activation. In addition, because of the spectral absorption of pyridoxal phosphate and the possible inhibition of the coupling enzyme used in the indicator reaction of the assay, higher concentrations of pyridoxal phosphate are not used in solution assays (see Hafkenscheid et al, "Influence of Pyridoxal-5'-phosphate on the Determination of the Alanine Aminotransferase and Aspartate Aminotransferase of Commercial Test Sera", J. Clin. Chem. Clin. Biochem., Vol. 17, pp. 219-223, (1979); Soo-se Chen et al, "Modification of Pig M.sub.4 Lactate Dehydrogenase by Pyridoxal 5'-Phosphate", Biochem J., Vol. 149, pp. 107-113, (1975) and "Reversible Modification of Pig Heart Mitochondrial Malate Dehydrogenase by Pyridoxal 5'-Phosphate", Biochem. J., Vol. 151, pp. 297-303 (1975)).

Detailed Description Text (2):

The present invention is described in relation to dry analytical elements for the determination of AST and ALT. It will be understood, however, that other enzymes which are activated by pyridoxal phosphate are quantitated by elements containing a high level of this activator. By choice of suitable detecting reagents, enzymes such as other transaminases, for example glutamate-cysteine transaminase, and other transaminases described in table 16-4 on page 684 of Mahler et al, Biological Chemistry, Harper and Row, 1966; isomerases, for example isomers for analine, glutamate, proline, lysine and serine and dicarboxylases, for example those described in Mahler et al, cited above, at page 685, are determined.

Other Reference Publication (4):

Hafkenschied et al. "Influence of Pyridoxal-5'-phosphate on the Determination of the Alanine Aminotransferase and Aspartate Aminotransferase of Commercial Test Sera", J. Clin. Chem. Clin. Biochem., vol. 17, pp. 219-223 (1979).

**WEST**

Generate Collection

L15: Entry 5 of 10

File: USPT

Feb 20, 1996

DOCUMENT-IDENTIFIER: US 5492821 A

TITLE: Stabilized polyacrylic saccharide protein conjugates

## BSPR:

The linker group includes all of the atoms between the polymer main chain (which is made of the atoms bonded to each other to form the length and longitudinal axis of the polymer) and the amino acid residue of the protein attached to the linker and will separate the main chain of the polymer from the protein by two or more carbon atoms and include at least three hydroxyl groups. In a very important aspect of the invention the linker group will be a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide.

## DEPR:

Enzymes, antibodies, and the like are complex proteins each with a specific sequence of amino acids. The structure of the protein is critical to the activity of the protein such as the catalytic activity of an enzyme and the capacity of an antibody to recognize ligands.

## DEPR:

"Protein" means proteins, including proteins modified to include additional amino groups such as lysine groups, polypeptides, enzymes, antibodies, and the like, which are composed of a specific sequence of amino acids.

## DEPR:

The linker group includes all atoms between the polymer main chain and the amino acid residue of the protein attached to the linker group and will separate the main chain of the polymer from the protein by two or more carbon atoms and have at least three hydroxyl groups. As used herein, a linker is a precursor to a linker group. In a very important aspect of the invention the linker group is a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide. A linker group which includes hydroxyl groups stabilizes the protein; and while there is not necessarily a defined limited as to the exact number of carbon atoms in a linker group, steric and kinetic considerations limit the size of the linker group to a total of about 60 carbon atoms.

## DEPR:

After a linker is coupled with a protein to form a linker group, the availability of an abundant number of hydroxyl groups on the linker group such as on a saccharide linker group provides enhancement of hydrophilic amino acid residue contacts.

## DEPR:

Enzymes which can be part of the polymer protein compounds of the invention include enzymes related to the production of fructose, such as glucose isomerase, which operates at 60.degree. to 65.degree. C., hydrolysis of starch by .alpha.-amylase, which occurs at 85.degree. to 110.degree. C. and resolution of D,L-amino acids by amino acetolase at 50.degree. C. Other enzymes which can be a part of the polymer protein compounds of the invention are nucleases, including endonucleases.

## DEPR:

Enzymes which can be stabilized as described here include oxidoreductases, transferases, dehydrogenases, transaminases, peptidases, hydrolases, isomerases, mutases and the like. In addition to enzymes already described, other enzymes which could be stabilized herein include but are not limited to malate dehydrogenase; creatine kinase (CK, CPK); alkaline phosphatase (AP, ALP, ALK-phos); aspartate aminotransferase (AST, OT, GOT); alanine aminotransferase (ALT, PT, SGPT); gamma glutamyl transpeptidase (gammaGT, gammaGTP); glutamic oxalacetic transaminase (SGOT); glutamic pyruvic transaminase (GTP); alpha amylase; beta amylase; lactate dehydrogenase (LD, LDH, lactic dehydrogenase); glucose-6-phosphate dehydrogenase (G6PDH); hexokinase (HK); glucose dehydrogenase; glucose oxidase; peroxidase (HRP, HPO, HRPO, PO); glycerol dehydrogenase; glutamate dehydrogenase, cholesterol oxidase; cholesterol esterase; lipase; uricase; urease; glycerol kinase; aldolases; synthetases; nucleases; polymerases; and the like.

## DEPR:

The polymer herein can also be used to synthesize peptide bonds. Typically, equimolar solutions containing modified amino acid(s) are dissolved in a solvent system containing an organic solvent such as dioxane or THF (tetrahydrofuran), or a tertiary amine such as (iPr).sub.2 NEt, to form an amino acid mixture. More preferably, the organic solvent is acetonitrile. The solutions dissolved in the solvent system are defined herein as amino acid acceptor and amino acid donor solutions. Examples of such solutions can be found in Table 2 of Example 10.

## DEPR:

A conjugated enzyme such as protease (the enzyme incorporated into one of the compounds of the invention) may be added to the amino acid mixture in an amount of about 10.sup.-6 to about 10.sup.-1 mole fraction relative to the amount of amino acid. The conjugated enzyme can be prepared as in Example 8. The reaction is performed at from about 0.degree. to 70.degree. C., at about 1 to about 24 hours. The enzyme polymer conjugate is removed from the formed peptide by filtering and the solvent is evaporated to obtain the resulting peptide.

## DETL:

## TABLE 2

Enzyme (Solvent, Time) Acceptor Donor Isolated (HPLC) Yield at 23.degree. C.  
Amino Acid Amino Acid Product (%)

CPC-CT	AcPhe--OEt	Ala--NH.sub.2	AcPheAla--NH.sub.2	97 (100)	(Dioxane, 12 h)
CPC-CT*	AcPhe--OEt	Ala--NH.sub.2	AcPheAla--NH.sub.2	94 (98)	(THF, 12 h)
CPC-BPN'	CbzLeuLeu--OMe	Leu--O.sup.t	Bu CbzLeuLeuLeu--O.sup.t	68 (95)	(Acetonitrile, 22 h)
CPC-BPN'	CbzLeuLeu--OMe	PheLeu--O.sup.t	Bu CbzLeuLeuPheLeu--O.sup.t	92 (98)	(Acetonitrile, 24 h)
CPC-BPN'	CbzValLeu--OMe	PheLeu--O.sup.t	Bu CbzValLeuPheLeu--O.sup.t	95 (98)	(Acetonitrile, 24 h)
CPC-BPN'	CbzValLeu--OMe	Ala--NH.sub.2	Ala--NH.sub.2	65 (95)	(Acetonitrile, 24 h)
CPC-BPN'	CbzLeuLeuAla--NH.sub.2	Ala--NH.sub.2	Ala--NH.sub.2	55 (90)	(Acetonitrile, 24 h)
CPC-T	CbzPhe--OH	Leu--OMe	CbzPheLeu--OMe	65 (95)	(Acetonitrile, 24 h)

50.degree. C.

\*at

## ORPL:

Margolin et al., Incorporation of D-Amino Acids into Peptides via Enzymatic Condensation in Organic Solvents, 109 J. Am. Chem. Soc. 7885-7887 (1987).

## ORPL:

West & Wong, Enzyme-Catalyzed Synthesis of Peptides Containing D-Amino Acids, J. Chem. Soc., Chem. Commun. 417-418 (1986).

**WEST**

Generate Collection

L15: Entry 4 of 10

File: USPT

Jun 17, 1997

DOCUMENT-IDENTIFIER: US 5639633 A

TITLE: Method for synthesizing peptides with saccharide linked enzyme polymer conjugates

## BSPR:

The linker group includes all of the atoms between the polymer main chain (which is made of the atoms bonded to each other to form the length and longitudinal axis of the polymer) and the amino acid residue of the protein attached to the linker and will separate the main chain of the polymer from the protein by two or more carbon atoms and include at least three hydroxyl groups. In a very important aspect of the invention the linker group will be a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide.

## DEPR:

Enzymes, antibodies, and the like are complex proteins each with a specific sequence of amino acids. The structure of the protein is critical to the activity of the protein such as the catalytic activity of an enzyme and the capacity of an antibody to recognize ligands.

## DEPR:

"Protein" means proteins, including proteins modified to include additional amino groups such as lysine groups, polypeptides, enzymes, antibodies, and the like, which are composed of a specific sequence of amino acids.

## DEPR:

The linker group includes all atoms between the polymer main chain and the amino acid residue of the protein attached to the linker group and will separate the main chain of the polymer from the protein by two or more carbon atoms and have at least three hydroxyl groups. As used herein, a linker is a precursor to a linker group. In a very important aspect of the invention the linker group is a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide. A linker group which includes hydroxyl groups stabilizes the protein; and while there is not necessarily a defined limited as to the exact number of carbon atoms in a linker group, steric and kinetic considerations limit the size of the linker group to a total of about 60 carbon atoms.

## DEPR:

After a linker is coupled with a protein to form a linker group, the availability of an abundant number of hydroxyl groups on the linker group such as on a saccharide linker group provides enhancement of hydrophilic amino acid residue contacts.

## DEPR:

Enzymes which can be part of the polymer protein compounds of the invention include enzymes related to the production of fructose, such as glucose isomerase, which operates at 60.degree. to 65.degree. C., hydrolysis of starch by .alpha.-amylase, which occurs at 85.degree. to 110.degree. C. and resolution of D,L-amino acids by amino acetolase at 50.degree. C. Other enzymes which can be a part of the polymer protein compounds of the invention are nucleases, including endonucleases.

## DEPR:

Enzymes which can be stabilized as described here include oxidoreductases, transferases, dehydrogenases, transaminases, peptidases, hydrolases, isomerases, mutases and the like. In addition to enzymes already described, other enzymes which could be stabilized herein include but are not limited to malate dehydrogenase; creatine kinase (CK, CPK); alkaline phosphatase (AP, ALP, ALK-phos); aspartate aminotransferase (AST, OT, GOT); alanine aminotransferase (ALT, PT, SGPT); gamma glutamyl transpeptidase (gammaGT, gammaGTP); glutamic oxalacetic transaminase (SGOT); glutamic pyruvic transaminase (GTP); alpha amylase; beta amylase; lactate dehydrogenase (LD, LDH, lactic dehydrogenase); glucose-6-phosphate dehydrogenase (G6PDH); hexokinase (HK); glucose dehydrogenase; glucose oxidase; peroxidase (HRP, HPO, HRPO, PO); glycerol dehydrogenase; glutamate dehydrogenase, cholesterol oxidase; cholesterol esterase; lipase; uricase; urease; glycerol kinase; aldolases; synthetases; nucleases; polymerases; and the like.

## DEPR:

The polymer herein can also be used to synthesize peptide bonds. Typically, equimolar solutions containing modified amino acid(s) are dissolved in a solvent system containing an organic solvent such as dioxane or THF (tetrahydrofuran), or a tertiary amine such as (iPr).sub.2 NEt, to form an amino acid mixture. More preferably, the organic solvent is acetonitrile. The solutions dissolved in the solvent system are defined herein as amino acid acceptor and amino acid donor solutions. Examples of such solutions can be found in Table 2 of Example 10.

## DEPR:

A conjugated enzyme such as protease (the enzyme incorporated into one of the compounds of the invention) may be added to the amino acid mixture in an amount of about 10.sup.-6 to about 10.sup.-1 mole fraction relative to the amount of amino acid. The conjugated enzyme can be prepared as in Example 8. The reaction is performed at from about 0.degree. to 70.degree. C., at about 1 to about 24 hours. The enzyme polymer conjugate is removed from the formed peptide by filtering and the solvent is evaporated to obtain the resulting peptide.

## DETL:

## TABLE 2

Enzyme (Solvent, Time) Acceptor Donor Isolated (HPLC) at 23.degree. C. Amino Acid Amino Acid Product Yield (%)

CPC-CT AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH.sub.2 97 (100) (Dioxane, 12 h)  
 CPC-CT\* AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH.sub.2 94 (98) (THF, 12h) CPC-BPN'  
 CbzLeuLeu-OMe Leu-O.sup.t Bu CbzLeuLeuLeu-O.sup.t Bu 68 (95) (Acetonitrile, 22 h)  
 CPC-BPN' CbzLeuLeu-OMe PheLeu-O.sup.t Bu CbzLeuLeuPheLeu-O.sup.t Bu 92 (98)  
 (Acetronitrile, 24 h) CPC-BPN' CbzValLeu-OMe PheLeu-O.sup.t Bu  
 CbzValLeuPheLeu-O.sup.t Bu 95 (98) (Acetronitrile, 24 h) CPC-BPN'  
 CbzValLeu-OMe Ala-NH.sub.2 CbzValLeuAla-NH.sub.2 65 (95) (Acetronitrile, 24 h)  
 CPC-BPN' CbzLeuLeu-OMe Ala-NH.sub.2 CbzLeuLeuAla-NH.sub.2 55 (90)  
 (Acetronitrile, 24 h) CPC-T CbzPhe-OH Leu-OMe CbzPheLeu-OMe 65 (95)  
 (Acetronitrile, 24 h)

\*at

50.degree. C.

## CLPR:

1. A method for making a peptide, the method comprising reacting at least two amino acids with a water soluble enzyme polymer conjugate to produce a peptide and recovering the peptide, the enzyme polymer conjugate comprising covalently bonded enzyme to a polymer, wherein the polymer is selected from the group consisting of a polyvinyl polymer, a polyester polymer, a polyamide polymer and mixtures thereof, the polymer having a main chain molecular weight of at least 5,000, and wherein three or more linker groups covalently bond both the enzyme and polymer and said linking groups have at least three hydroxyl groups and not more than 60 carbon atoms, each of said linker groups being selected



from the group consisting of ##STR26## wherein E---OH or --SH;

**WEST**

Generate Collection

L15: Entry 2 of 10

File: USPT

Apr 7, 1998

DOCUMENT-IDENTIFIER: US 5736625 A

TITLE: Method for stabilizing proteins with saccharide linked protein polymer conjugates

## BSPR:

The linker group includes all of the atoms between the polymer main chain (which is made of the atoms bonded to each other to form the length and longitudinal axis of the polymer) and the amino acid residue of the protein attached to the linker and will separate the main chain of the polymer from the protein by two or more carbon atoms and include at least three hydroxyl groups. In a very important aspect of the invention the linker group will be a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide.

## DEPR:

Enzymes, antibodies, and the like are complex proteins each with a specific sequence of amino acids. The structure of the protein is critical to the activity of the protein such as the catalytic activity of an enzyme and the capacity of an antibody to recognize ligands.

## DEPR:

"Protein" means proteins, including proteins modified to include additional amino groups such as lysine groups, polypeptides, enzymes, antibodies, and the like, which are composed of a specific sequence of amino acids.

## DEPR:

The linker group includes all atoms between the polymer main chain and the amino acid residue of the protein attached to the linker group and will separate the main chain of the polymer from the protein by two or more carbon atoms and have at least three hydroxyl groups. As used herein, a linker is a precursor to a linker group. In a very important aspect of the invention the linker group is a saccharide group which includes the residue of a monosaccharide, a disaccharide and a trisaccharide. A linker group which includes hydroxyl groups stabilizes the protein; and while there is not necessarily a defined limit as to the exact number of carbon atoms in a linker group, steric and kinetic considerations limit the size of the linker group to a total of about 60 carbon atoms.

## DEPR:

After a linker is coupled with a protein to form a linker group, the availability of an abundant number of hydroxyl groups on the linker group such as on a saccharide linker group provides enhancement of hydrophilic amino acid residue contacts.

## DEPR:

Enzymes which can be part of the polymer protein compounds of the invention include enzymes related to the production of fructose, such as glucose isomerase, which operates at 60.degree. to 65.degree. C., hydrolysis of starch by .alpha.-amylase, which occurs at 85.degree. to 110.degree. C. and resolution of D,L-amino acids by amino acetolase at 50.degree. C. Other enzymes which can be a part of the polymer protein compounds of the invention are nucleases, including endonucleases.

## DEPR:

Enzymes which can be stabilized as described here include oxidoreductases, transferases, dehydrogenases, transaminases, peptidases, hydrolaser, isomerases, mutases and the like. In addition to enzymes already described, other enzymes which could be stabilized herein include but are not limited to malate dehydrogenase; creatine kinase (CK, CPK); alkaline phosphatase (AP, ALP, ALK-phos); aspartate aminotransferase (AST, OT, GOT); alanine aminotransferase (ALT, PT, SGPT); gamma glutamyl transpeptidase (gammaGT, gammaGTP); glutamic oxalacetic transaminase (SGOT); glutamic pyruvic transaminase (GTP); alpha amylase; beta amylase; lactate dehydrogenase (LD, LDH, lactic dehydrogenase); glucose-6-phosphate dehydrogenase (G6PDH); hexokinase (HK); glucose dehydrogenase; glucose oxidase; peroxidase (HRP, HPO, HRPO, PO); glycerol dehydrogenase; glutamate dehydrogenase, cholesterol oxidase; cholesterol esterase; lipase; uricase; urease; glycerol kinase; aldolases; synthetases; nucleases; polymerases; and the like.

## DEPR:

The polymer herein can also be used to synthesize peptide bonds. Typically, equimolar solutions containing modified amino acid(s) are dissolved in a solvent system containing an organic solvent such as dioxane or THF (tetrahydrofuran), or a tertiary amine such as (iPr).sub.2 NEt, to form an amino acid mixture. More preferably, the organic solvent is acetonitrile. The solutions dissolved in the solvent system are defined herein as amino acid acceptor and amino acid donor solutions. Examples of such solutions can be found in Table 2 of Example 10.

## DEPR:

A conjugated enzyme such as protease (the enzyme incorporated into one of the compounds of the invention) may be added to the amino acid mixture in an amount of about 10.sup.-6 to about 10.sup.-1 mole fraction relative to the amount of amino acid. The conjugated enzyme can be prepared as in Example 8. The reaction is performed at from about 0.degree. to 70.degree. C., at about 1 to about 24 hours. The enzyme polymer conjugate is removed from the formed peptide by filtering and the solvent is evaporated to obtain the resulting peptide.

## DETL:

## TABLE 2

Enzyme (Solvent, Time) Acceptor Donor Isolated (HPLC) at 23.degree. C. Amino Acid Amino Acid Product Yield (%)

CPC-CT AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH.sub.2 97(100) (Dioxane, 12 h) CPC-CT  
 \* AcPhe-OEt Ala-NH.sub.2 AcPheAla-NH<sub>2</sub> 94(98) (THF, 12 h) CPC-BPN'  
 CbzLeuLeu-OMe Leu-O.sup.t Bu CbzLeuLeuLeu-O.sup.t Bu 68(95) (Acetonitrile, 22 h) CPC-BPN' CbzLeuLeu-OMe PheLeu-O.sup.t Bu CbzLeuLeuPheLeu-O.sup.t Bu 92(98) (Acetronitrile, 24 h) CPC-BPN' CbzValLeu-OMe PheLeu-O.sup.t Bu CbzValLeuPheLeu-O.sup.t Bu 95(98) (Acetronitrile, 24 h) CPC-BPN' CbzValLeu-OMe Ala-NH.sub.2 CbzValLeuAla-NH.sub.2 65(95) (Acetronitrile, 24 h) CPC-BPN' CbzLeuLeu-OMe Ala-NH.sub.2 CbzLeuLeuAla-NH.sub.2 55(90) (Acetronitrile, 24 h) CPC-T CbzPhe-OH Leu-OMe CbzPheLeu-OMe 65(95) (Acetronitrile, 24 h)

at 50.degree. C.

\*

**WEST**

Generate Collection

Print

L8: Entry 51 of 71

File: USPT

Sep 8, 1998

DOCUMENT-IDENTIFIER: US 5804402 A

TITLE: Reagent

Abstract Text (1):

This invention relates to a reagent for enzymatic determination of an analyte concentration in a patient wherein the degree of oxidation of a coenzyme is measured, characterized in that said reagent is stabilized against oxidation by a coenzyme reduction system comprising an enzyme and substrate pair selected so as to enable continuous regeneration of said coenzyme throughout storage of said reagent. Also disclosed is an improvement in an enzymatic method of determination of an analyte concentration in a sample body fluid wherein the degree of oxidation of a coenzyme is measured, the improvement comprising stabilizing a reagent comprising said coenzyme against oxidation by a coenzyme reduction system comprising an enzyme and substrate pair selected so as to enable continuous regeneration of said coenzyme throughout storage of said reagent. Also disclosed are reagents for the determination of aspartate aminotransferase, alanine aminotransferase, ammonia and urea.

Brief Summary Text (7):

The indirect quantification of enzymes, in particular, the transaminases, aspartate aminotransferase and alanine aminotransferase in sample body fluids may involve contrasting a sample "blank" against a sample in which the enzymatic conversion of an analyte associated with the enzyme of interest has taken place.

Brief Summary Paragraph Table (3):

	Substrate Relative Activity
	<u>L-valine</u> 74% L-isoleucine 58% L-norvaline 41%
L-norleucine 10% L-methionine 0.6% L-cysteine 0.3%	

**WEST**

Generate Collection

Print

L8: Entry 50 of 71

File: USPT

Sep 29, 1998

DOCUMENT-IDENTIFIER: US 5814473 A

**\*\* See image for Certificate of Correction \*\***

TITLE: Transaminases and aminotransferases

Brief Summary Text (5):

The measurement of alanine aminotransferase and aspartate aminotransferase levels in blood serum is an important diagnostic procedure in medicine, used as an indicator of heart damage and to monitor recovery from the damage.

Detailed Description Text (68):

Transaminases are highly stereoselective, and most use L-amino acids as substrates. Using the approach disclosed in a commonly assigned, copending provisional application Ser. No. 60/008,316, filed on Dec. 7, 1995 and entitled "Combinatorial Enzyme Development," the disclosure of which is incorporated herein by reference in its entirety, one can convert the transaminases of the invention to use D-amino acids as substrates. Such conversion makes possible a broader array of transaminase applications. For instance, D-valine can be used in the manufacture of synthetic pyrethroids. D-phenylglycine and its derivatives can be useful as components of .beta.-lactam antibiotics. Further, the thermostable transaminases have superior stability at higher temperatures and in organic solvents. Thus, they are better suited to utilize either L- and/or D-amino acids for production of optically pure chiral compounds used in pharmaceutical, agricultural, and other chemical manufactures.

**WEST**

Generate Collection

Print

L8: Entry 44 of 71

File: USPT

Oct 5, 1999

DOCUMENT-IDENTIFIER: US 5962283 A

TITLE: Transminases and amnottransferases

Brief Summary Text (5):

The measurement of alanine aminotransferase and aspartate aminotransferase levels in blood serum is an important diagnostic procedure in medicine, used as an indicator of heart damage and to monitor recovery from the damage.

Brief Summary Text (81):

Transaminases are highly stereoselective, and most use L-amino acids as substrates. Using the approach disclosed in a commonly assigned, copending provisional application Ser. No. 60/008,316, filed on Dec. 7, 1995 and entitled "Combinatorial Enzyme Development," the disclosure of which is incorporated herein by reference in its entirety, one can convert the transaminases of the invention to use D-amino acids as substrates. Such conversion makes possible a broader array of transaminase applications. For instance, D-valine can be used in the manufacture of synthetic pyrethroids. D-phenylglycine and its derivatives can be useful as components of .beta.-lactam antibiotics. Further, the thermostable transaminases have superior stability at higher temperatures and in organic solvents. Thus, they are better suited to utilize either L- and/or D-amino acids for production of optically pure chiral compounds used in pharmaceutical, agricultural, and other chemical manufactures.

**WEST**

## Freeform Search

**Database:**

US Patents Full-Text Database  
US Pre-Grant Publication Full-Text Database  
JPO Abstracts Database  
EPO Abstracts Database  
Derwent World Patents Index  
IBM Technical Disclosure Bulletins

**Term:**

114 and 113

**Display:**

10

**Documents in Display Format:**

CIT

**Starting with Number**

1

**Generate:** ☐ Hit List ☒ Hit Count ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

### Search History

**Today's Date:** 4/26/2001

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l14 and l13	10	<u>L15</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l2 or (amino acid )	145677	<u>L14</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l5 same l1	24	<u>L13</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l5 and l1	390	<u>L12</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l9 not l10	8	<u>L11</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l5 and l9	11	<u>L10</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l7 same l2	19	<u>L9</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l7 and l5	645	<u>L8</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	aminotransferase	1166	<u>L7</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l1 same l2	5	<u>L6</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	stabiliz\$5 or stabl\$5	1264710	<u>L5</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	stabiliz\$5	505158	<u>L4</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	l1 and l2	108	<u>L3</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	valine or proline	27431	<u>L2</u>
USPT,PGPB,JPAB,EPAB,DWPI,TDBD	(aspartate aminotransferase) or (alanine aminotransferase)	674	<u>L1</u>



Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:sssptal815mxw  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 3 Feb 06 Engineering Information Encompass files have new names  
NEWS 4 Feb 16 TOXLINE no longer being updated  
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure  
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA

NEWS EXPRESS April 18 CURRENT WINDOWS VERSION IS V6.0,  
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),  
AND CURRENT DISCOVER FILE IS DATED 04/06

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:20:46 ON 26 APR 2001

=> index bioscience

FILE 'DRUGMONOG' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.15	0.15

FULL ESTIMATED COST

INDEX 'ADISALERTS, ADISINSIGHT, AGRICOLA, ANABSTR, AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DRUGB, DRUGLAUNCH, DRUGMONOG2, DRUGNL, ...' ENTERED AT 12:21:11 ON 26 APR 2001

59 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view search error messages that display as 0\* with SET DETAIL OFF.

=> file reg

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.45	0.60

FULL ESTIMATED COST

FILE 'REGISTRY' ENTERED AT 12:21:39 ON 26 APR 2001  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 25 APR 2001 HIGHEST RN 332836-91-6  
DICTIONARY FILE UPDATES: 25 APR 2001 HIGHEST RN 332836-91-6

TSCA INFORMATION NOW CURRENT THROUGH January 11, 2001

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Structure search limits have been increased. See HELP SLIMIT  
for details.

=> s aspartate aminotransferase/cn  
L1 1 ASPARTATE AMINOTRANSFERASE/CN

=> d

L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2001 ACS  
RN 9000-97-9 REGISTRY  
CN Aminotransferase, aspartate (9CI) (CA INDEX NAME)  
OTHER NAMES:  
CN 2-Oxoglutarate-glutamate aminotransferase  
CN Aspartate .alpha.-ketoglutarate transaminase  
CN **Aspartate aminotransferase**  
CN Aspartate-2-oxoglutarate transaminase  
CN Aspartic acid aminotransferase  
CN Aspartic aminotransferase  
CN Aspartyl aminotransferase  
CN AST  
CN E.C. 2.6.1.1  
CN Glutamate 2-oxoglutarate transaminase  
CN Glutamate-oxalacetate aminotransferase  
CN Glutamate-oxalate transaminase  
CN Glutamate-oxaloacetate transaminase  
CN Glutamic-aspartic aminotransferase  
CN Glutamic-aspartic transaminase  
CN Glutamic-oxalacetic transaminase  
CN Glutamic-oxalic transaminase  
CN GOT  
CN GOT (enzyme)  
CN L-Aspartate aminotransferase  
CN L-Aspartate transaminase  
CN L-Aspartate-.alpha.-ketoglutarate transaminase  
CN L-Aspartate-2-ketoglutarate aminotransferase  
CN L-Aspartate-2-oxoglutarate aminotransferase  
CN L-Aspartate-2-oxoglutarate-transaminase  
CN L-Aspartic aminotransferase  
CN Oxalacetate-aspartate aminotransferase  
CN Oxaloacetate transferase  
DR 9013-64-3, 9014-29-3, 9016-19-7, 9036-26-4, 9061-83-0, 61461-53-8,  
139074-52-5  
MF Unspecified  
CI MAN  
LC STN Files: AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA,  
CAPLUS, CASREACT, CHEMCATS, CHEMLIST, CIN, CSCHEM, EMBASE, IFICDB,

IFIPAT, IFIUDB, IPA, MEDLINE, MSDS-OHS, NAPRALERT, NIOSHTIC, PROMT,  
TOXLINE, TOXLIT, USPATFULL  
Other Sources: EINECS\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

14782 REFERENCES IN FILE CA (1967 TO DATE)

85 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

14805 REFERENCES IN FILE CAPLUS (1967 TO DATE)

=> s alanine aminotransferase/cn  
L2 1 ALANINE AMINOTRANSFERASE/CN

=> d

L2 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2001 ACS

RN 9000-86-6 REGISTRY

CN Aminotransferase, alanine (8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN **Alanine aminotransferase**

CN Alanine transaminase

CN Alanine-.alpha.-ketoglutarate aminotransferase

CN Alanine-oxaloacetate aminotransferase

CN Alanine-pyruvate aminotransferase

CN ALT

CN E.C. 2.6.1.2

CN Glutamic acid-pyruvic acid transaminase

CN Glutamic-alanine transaminase

CN Glutamic-pyruvic aminotransferase

CN Glutamic-pyruvic transaminase

CN GPT

CN L-Alanine aminotransferase

CN L-Alanine transaminase

CN L-Alanine-.alpha.-ketoglutarate aminotransferase

CN Pyruvate transaminase

CN Pyruvate-alanine aminotransferase

CN Pyruvate-glutamate transaminase

CN SGPT

DR 9013-65-4, 9014-30-6, 9048-64-0, 139074-53-6

MF Unspecified

CI MAN

LC STN Files: AGRICOLA, ANABSTR, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA,  
CAPLUS, CASREACT, CHEMCATS, CHEMLIST, CIN, CSCHEM, CSNB, EMBASE,

IFICDB,

IFIPAT, IFIUDB, MEDLINE, MSDS-OHS, NAPRALERT, NIOSHTIC, PROMT, TOXLINE,  
TOXLIT, USPATFULL

Other Sources: EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

10974 REFERENCES IN FILE CA (1967 TO DATE)

14 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

10995 REFERENCES IN FILE CAPLUS (1967 TO DATE)

=> index bioscience

FILE 'DRUGMONOG' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION

FULL ESTIMATED COST

11.22

11.82

INDEX 'ADISALERTS, ADISINSIGHT, AGRICOLA, ANABSTR, AQUASCI, BIOBUSINESS,  
BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT,  
CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE,  
DRUGB, DRUGLAUNCH, DRUGMONOG2, DRUGNL, ...' ENTERED AT 12:22:47 ON 26  
APR 2001

59 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view  
search error messages that display as 0\* with SET DETAIL OFF.

=> s 11 or 12

0\* FILE ADISALERTS  
1000 FILE AGRICOLA  
140 FILE ANABSTR  
0\* FILE AQUASCI  
304 FILE BIOBUSINESS  
0\* FILE BIOCOMMERCE  
7779 FILE BIOSIS  
2676 FILE BIOTECHNO  
0\* FILE CABA  
0\* FILE CAPLUS  
0\* FILE CEABA-VTB  
8 FILE CIN  
0\* FILE CONFSCI  
0\* FILE CROPB  
0\* FILE CROPU  
0\* FILE DDFB  
0\* FILE DDFU  
0\* FILE DGENE  
0\* FILE DRUGB  
0\* FILE DRUGU  
0\* FILE EMBAL  
0\* FILE ESBIODBASE  
0\* FILE FOMAD  
0\* FILE FOREGE  
0\* FILE FROSTI  
0\* FILE GENBANK  
0\* FILE HEALSAFE  
0\* FILE IFIPAT  
0\* FILE KOSMET  
0\* FILE LIFESCI  
0\* FILE MEDICONF  
1264 FILE MEDLINE  
45 FILES SEARCHED...  
0\* FILE NTIS  
0\* FILE OCEAN  
0\* FILE PASCAL  
0\* FILE PHIC  
0\* FILE PHIN  
119 FILE PROMT  
0\* FILE SCISEARCH  
0\* FILE USPATFULL

8 FILES HAVE ONE OR MORE ANSWERS, 59 FILES SEARCHED IN STNINDEX

L3 QUE L1 OR L2

=> s valine? or proline?

102	FILE ADISALERTS
62	FILE ADISINSIGHT
3361	FILE AGRICOLA
988	FILE ANABSTR
923	FILE AQUASCI
1357	FILE BIOBUSINESS
47	FILE BIOCOMMERCE
39518	FILE BIOSIS
2172	FILE BIOTECHABS
2172	FILE BIOTECHDS
11612	FILE BIOTECHNO
10774	FILE CABA
4033	FILE CANCERLIT
78892	FILE CAPLUS
453	FILE CEABA-VTB
55	FILE CEN
47	FILE CIN
483	FILE CONFSCI
149	FILE CROPB
393	FILE CROPU
2061	FILE DDFB
1688	FILE DDFU
3445	FILE DGENE
2061	FILE DRUGB
12	FILE DRUGLAUNCH
984	FILE DRUGMONOG2
35	FILE DRUGNL
2480	FILE DRUGU
59	FILE DRUGUPDATES
200	FILE EMBAL
27954	FILE EMBASE
7795	FILE ESBIODBASE
1	FILE FOMAD
47	FILE FOREGE
761	FILE FROSTI
35 FILES SEARCHED...	
2685	FILE FSTA
5967	FILE GENBANK
47	FILE HEALSAFE
3325	FILE IFIPAT
1975	FILE JICST-EPLUS
48	FILE KOSMET
9234	FILE LIFESCI
35095	FILE MEDLINE
331	FILE NIOSHTIC
304	FILE NTIS
253	FILE OCEAN
9871	FILE PASCAL
112	FILE PHAR
1	FILE PHIC
49	FILE PHIN
1646	FILE PROMT
22030	FILE SCISEARCH
117	FILE SYNTHLINE
6734	FILE TOXLINE
9935	FILE TOXLIT
20991	FILE USPATFULL
4548	FILE WPIDS
4548	FILE WPINDEX

58 FILES HAVE ONE OR MORE ANSWERS, 59 FILES SEARCHED IN STNINDEX

L4 QUE VALINE? OR PROLINE?

=> s (aspartate aminotransferase? ) or (alanine aminotransferase?)

1067 FILE ADISALERTS  
39 FILE ADISINSIGHT  
1059 FILE AGRICOLA  
261 FILE ANABSTR  
279 FILE AQUASCI  
316 FILE BIOBUSINESS  
23 FILE BIOCOMMERCE  
12011 FILE BIOSIS  
113 FILE BIOTECHABS  
113 FILE BIOTECHDS  
3675 FILE BIOTECHNO  
5911 FILE CABA  
1495 FILE CANCERLIT  
12794 FILE CAPLUS  
39 FILE CEABA-VTB  
8 FILE CIN  
116 FILE CONFSCI  
19 FILE CROPB  
321 FILE CROPU  
164 FILE DDFB  
2190 FILE DDFU  
861 FILE DGENE  
164 FILE DRUGB  
10 FILE DRUGNL  
2974 FILE DRUGU  
13 FILE DRUGUPDATES  
79 FILE EMBAL  
20823 FILE EMBASE  
2182 FILE ESBIODBASE

32 FILES SEARCHED...

58 FILE FROSTI  
97 FILE FSTA  
1728 FILE GENBANK  
98 FILE HEALSAFE  
52 FILE IFIPAT  
566 FILE JICST-EPLUS  
6 FILE KOSMET  
2184 FILE LIFESCI  
9483 FILE MEDLINE  
546 FILE NIOSHTIC  
93 FILE NTIS  
72 FILE OCEAN  
3478 FILE PASCAL  
7 FILE PHAR  
61 FILE PHIN  
412 FILE PROMT  
7042 FILE SCISEARCH  
3734 FILE TOXLINE  
3522 FILE TOXLIT  
531 FILE USPATFULL  
101 FILE WPIDS  
101 FILE WPINDEX

51 FILES HAVE ONE OR MORE ANSWERS, 59 FILES SEARCHED IN STNINDEX

Q

N

Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:sssptal815mxw  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 3 Feb 06 Engineering Information Encompass files have new names  
NEWS 4 Feb 16 TOXLINE no longer being updated  
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure  
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA

NEWS EXPRESS April 18 CURRENT WINDOWS VERSION IS V6.0,  
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),  
AND CURRENT DISCOVER FILE IS DATED 04/06  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:56:36 ON 26 APR 2001

=> file ca, biosis, medline  
COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
0.15	0.15

FILE 'CA' ENTERED AT 12:57:06 ON 26 APR 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 12:57:06 ON 26 APR 2001  
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'MEDLINE' ENTERED AT 12:57:06 ON 26 APR 2001

=> s (aspartate aminotransferase?) or (alanine aminotransferase?)  
L1 34077 (ASPARTATE AMINOTRANSFERASE?) OR (ALANINE AMINOTRANSFERASE?)

=> s valine or proline  
L2 151467 VALINE OR PROLINE



=> s 11 (p) 12  
L3 238 L1 (P) L2

=> s stab?  
L4 1677944 STAB?

=> s 13 and 14  
L5 16 L3 AND L4

=> dup re, 15  
'RE' IS NOT VALID HERE  
Enter "REMOVE" to identify and remove duplicate answers.  
Enter "IDENTIFY" to identify duplicate answers in the answer set.  
Enter "ONLY" to identify and create an answer set containing only duplicate records.  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):dup rem 15  
'DUP REM L20' IS NOT VALID HERE  
Enter "REMOVE" to identify and remove duplicate answers.  
Enter "IDENTIFY" to identify duplicate answers in the answer set.  
Enter "ONLY" to identify and create an answer set containing only duplicate records.  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):.  
Enter "REMOVE" to identify and remove duplicate answers.  
Enter "IDENTIFY" to identify duplicate answers in the answer set.  
Enter "ONLY" to identify and create an answer set containing only duplicate records.  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):d 1-16 ab,bib  
'D 1-16 AB,BIB' IS NOT VALID HERE  
Enter "REMOVE" to identify and remove duplicate answers.  
Enter "IDENTIFY" to identify duplicate answers in the answer set.  
Enter "ONLY" to identify and create an answer set containing only duplicate records.  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):.  
Enter "REMOVE" to identify and remove duplicate answers.  
Enter "IDENTIFY" to identify duplicate answers in the answer set.  
Enter "ONLY" to identify and create an answer set containing only duplicate records.  
ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove  
PROCESSING COMPLETED FOR L5  
L6 10 DUP REMOVE L5 (6 DUPLICATES REMOVED)

=> d 1-10 ab,bib

L6 ANSWER 1 OF 10 CA COPYRIGHT 2001 ACS  
AB **Aspartate aminotransferase** or **alanine aminotransferase** in liq. media such as serum or buffered soln. is **stabilized** by adding **valine**, **proline**, or both into the media to assure the accuracy of clin. studies. The enzymes can be further **stabilized** by adding a sol. protein such as albumin or gelatin into the media.  
AN 131:319672 CA  
TI **Stabilization** of aminotransferase using amino acids  
IN Baba, Toshiyuki; Tabata, Hiromasa; Nagamatsu, Katashi; Watazu, Yoshifumi; Aoki, Ryoji  
PA International Reagents Corporation, Japan; Asahi Chemical Industry Co., Ltd.  
SO PCT Int. Appl., 30 pp.  
CODEN: PIXXD2  
DT Patent  
LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9955850	A1	19991104	WO 1999-JP2205	19990423
	W: CN, JP, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
PRAI	JP 1998-131159		19980424		

RE.CNT 9

RE

- (2) Beckman Instruments Inc; EP 16573 A CA
  - (3) Beckman Instruments Inc; US 4325832 A CA
  - (4) Beckman Instruments Inc; JP 55-141194 A 1980 CA
  - (5) Iwan Endore Modorobich; US 4652524 A CA
  - (6) Iwan Endore Modorobich; EP 49475 A CA
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 10 CA COPYRIGHT 2001 ACS

DUPLICATE 1

AB To elucidate the role of the two conserved **cis-proline** residues of **aspartate aminotransferase** (AspAT), one double and two single mutants of the enzyme from *Escherichia coli* (EcAspAT) were prep.d.: P138A, P195A and P138A/P195A in which the two prolines were replaced by alanine. The crystal structures of P195A and P138A/P195A have been detd. at 2.3-2.1 .ANG. resolu. The wild-type geometry, including the cis conformation of the 194-195 peptide bond is retained upon substitution of **proline** 195 by alanine, whereas the trans conformation is adopted at the 137-138 peptide bond. Quite surprisingly, the replacement of each of the two prolines by alanine does not significantly affect either the activity or the **stability** of the protein. All the three mutants follow the same pathway as the wild type for unfolding equil. induced by guanidine hydrochloride [Herold, M., and Kirschner, K. (1990) *Biochem.* 29, 1907-1913]. The kinetics of renaturation of P195A, where the alanine retains the wild-type cis conformation, is faster than wild type, whereas renaturation of P138A, which adopts the trans conformation, is slower. We conclude that **cis-prolines** seem to have been retained throughout the evolution of **aspartate aminotransferase** to possibly play a subtle role in directing the traffic of intermediates toward the unique structure of the native state, rather than to respond to the needs for a specific catalytic or functional role.

AN 130:164805 CA

TI Functional and Structural Analysis of **cis-Proline** Mutants of *Escherichia coli* **Aspartate Aminotransferase**

AU Birolo, Leila; Malashkevich, Vladimir N.; Capitani, Guido; De Luca, Fabio;

CS Moretta, Alma; Jansonius, Johan N.; Marino, Gennaro  
Dipartimento di Chimica Organica e Biologica, Universita Federico II, Naples, I-80134, Italy

SO *Biochemistry* (1999), 38(3), 905-913  
CODEN: BICHAW; ISSN: 0006-2960

PB American Chemical Society

DT Journal

LA English

RE.CNT 45

RE

(1) Birolo, L; *Eur J Biochem* 1995, V232, P859 CA

- (3) Garnier, A; Eur J Biochem 1993, V216, P763 CA  
 (4) Gloss, L; Biochemistry 1992, V31, P32 CA  
 (5) Goldberg, J; Biochemistry 1991, V30, P305 CA  
 (6) Herold, M; Biochemistry 1990, V29, P1907 CA  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 3 OF 10 CA COPYRIGHT 2001 ACS  
 AB The adiabatic compressibility (.hivin..beta.s) was detd., by the precise  
 sound velocity and d. measurements, for a series of single amino acid  
 substituted mutant enzymes of Escherichia coli dihydrofolate reductase  
 (DHFR) and **aspartate aminotransferase** (AspAT).  
 Interestingly, the .hivin..beta.s values of both DHFR and AspAT were  
 influenced markedly by the mutations at glycine-121 and **valine**  
 -39, resp., in which the magnitude of the change was proportional to the  
 enzyme activity. This result demonstrates that the local change of the  
 primary structure plays an important role in at. packing and protein  
 dynamics, which leads to the modified **stability** and enzymic  
 function. This is the first report on the compressibility of mutant  
 proteins.  
 AN 124:224204 CA  
 TI A large compressibility change of protein induced by a single amino acid  
 substitution  
 AU Gekko, Kunihiro; Tamura, Youjiro; Ohmae, Eiji; Hayashi, Hideyuki;  
 Kagamiyama, Hiroyuki; Ueno, Hiroshi  
 CS Fac. Science, Hiroshima Univ., Higashi-Hiroshima, 739, Japan  
 SO Protein Sci. (1996), 5(3), 542-5  
 CODEN: PRCIEI; ISSN: 0961-8368  
 DT Journal  
 LA English

L6 ANSWER 4 OF 10 CA COPYRIGHT 2001 ACS  
 AB A notable feature of porcine cytosolic aspartate aminotransferase is the  
 closure of the active site cleft by a mobile amino-terminal segment  
 (residues 15-40) upon binding substrate. The functional roles of Val17  
 and Phe18, residues that are part of the mobile loop, have been studied  
 in the site-directed mutants in which the size and hydrophobic nature of  
 these residues have been changed. Absorption, CD spectra, susceptibility  
 to protease 401, and thermal **stability** did not differ  
 appreciably between wild type and mutant enzymes. In the overall  
 transamination between aspartate and 2-oxoglutarate, V17A represented a  
 typical Km mutant while V17I retained the substrate binding affinity  
 fairly well. In contrast, replacement of Phe18 by Ala resulted in a  
 large decrease in both catalytic rate and binding affinity for substrates.  
 F18W, F18Y, and F18H showed a moderate decrease in kcat and a  
 considerable increase in Km values. Single-turnover reactions with four individual  
 substrates yielded analogous results to those obtained for the overall  
 reaction and, in addn., revealed that k/Kd values of mutants F18A and  
 F18H were over 10 times lower for C5 substrates (glutamate and 2-oxoglutarate)  
 than those for C4 substrates (aspartate and oxalacetate). All mutant  
 enzymes showed variously increased Kd values for substrate analogs such  
 as 2-methylaspartate, succinate, and glutarate. 1H NMR observations of  
 F18H, in which His18 served as a built-in probe, were in accord with the  
 behavior that would be expected from the conformational transition. The  
 authors conclude that, although Val17 and Phe18 may not be essential for

- catalysis, the presence of a bulky residue of appropriate size at each position is crit. for productive binding of substrate.
- AN 121:173879 CA
- TI Functional role of the amino-terminal mobile segment in catalysis by porcine cytosolic aspartate aminotransferase. Critical importance of
- Vall17 and Phe18 for productive binding of substrates
- AU Nishimura, Kosuke; Higaki, Tsuyoshi; Okamura, Hitoshi; Tanase, Sumio
- CS Sch. Med., Kumamoto Univ., Kumamoto, 860, Japan
- SO J. Biol. Chem. (1994), 269(40), 24712-18
- CODEN: JBCHA3; ISSN: 0021-9258
- DT Journal
- LA English
- L6 ANSWER 5 OF 10 CA COPYRIGHT 2001 ACS DUPLICATE 3
- AB The retina of honeybee drone is a nervous tissue with a crystal-like structure in which glial cells and photoreceptor neurons constitute 2 distinct metabolic compartments. The phosphorylation of glucose and its subsequent incorporation into glycogen occur in glia, whereas O2 consumption (QO2) occurs in the photoreceptors. Exptl. evidence showed that glia phosphorylate glucose and supply the photoreceptors with metabolic substrates. The authors aimed to identify these transferred substrates. Using ion-exchange and reversed-phase HPLC and gas chromatog.-mass spectrometry, the authors demonstrated that >50% of 14C(U)-glucose entering the glia is transformed to alanine by transamination of pyruvate and glutamate. In the absence of extracellular glucose, glycogen is used to make alanine; thus, its pool size in isolated retinas is maintained **stable** or even increased. The authors' model proposes that the formation of alanine occurs in the glia, thereby maintaining the redox potential of this cell and contributing to NH3 homeostasis. Alanine is released into the extracellular space and is then transported into photoreceptors using an Na+-dependent transport system. Purified suspensions of photoreceptors have similar **alanine aminotransferase** activity as glial cells and transform 14C-alanine to glutamate, aspartate, and CO2. Therefore, the alanine entering photoreceptors is transaminated to pyruvate, which in turn enters the Krebs cycle. **Proline** also supplies the Krebs cycle by making glutamate and, in turn, the intermediate .alpha.-ketoglutarate. Light stimulation caused a 200% increase of QO2 and a 50% decrease of **proline** and of glutamate. Also, the prodn. of 14CO2 from 14C-**proline** was increased. The use of these amino acids would sustain about half of the light-induced .DELTA.QO2, the other half being sustained by glycogen via alanine formation. The use of **proline** meets a necessary anaplerotic function in the Krebs cycle, but implies high NH3 prodn. The results showed that alanine formation fixes NH3 at a rate exceeding glutamine formation. This is consistent with the rise of a glial pool of alanine upon photostimulation. In conclusion, the results strongly support a nutritive function for glia.
- AN 121:31478 CA
- TI Glial cells transform glucose to alanine, which fuels the neurons in the honeybee retina
- AU Tsacopoulos, M.; Veuthey, A. L.; Saravelos, S. G.; Perrottet, P.; Tsoupras, G.
- CS Sch. Med., Univ. Geneva, Geneva, 1211, Switz.
- SO J. Neurosci. (1994), 14(3, Pt. 1), 1339-51
- CODEN: JNRSDS; ISSN: 0270-6474

DT Journal  
LA English

L6 ANSWER 6 OF 10 MEDLINE

AB The functional roles of Val37 and Gly38 in porcine cytosolic aspartate aminotransferase have been studied in the site-directed mutants V37A, G38A, and G38S where the size and hydrophobic character of these residues has been altered. Previous x-ray studies have shown that Val37 and Gly38, which are part of a flexible loop, interact directly with bound substrate.

From x-ray and solution experiments we find that the V37A, G38A, and G38S mutations do not cause significant perturbations to the unliganded enzyme.

Replacing Val37 with a less bulky alanine residue does not affect the maximal catalytic rate (kcat), but it does increase significantly the Michaelis constants for substrates in the overall transamination reaction between aspartate and 2-oxoglutarate. On the other hand, replacing Gly38 with alanine or serine results in striking decreases in kcat to 5 and 0.6%, respectively, of the value observed for the wild-type enzyme, as well as in considerable increases in Km values. Consequently, the catalytic competence, kcat/Km, decreases by 3 orders of magnitude for

G38A

and by 4 orders of magnitude for G38S. Single turnover reactions of G38A and G38S with four individual substrates (aspartate, glutamate, oxalacetate, and 2-oxoglutarate) are characterized by kinetic parameters that are largely consistent with those of the overall reaction. In addition, the mutations at position 38 impair more seriously the catalytic

competence of the enzyme toward C5-substrates than toward C4-substrates. We conclude that Gly38 is probably required for proper function of the enzyme because it permits a high level of flexibility for the 36-39 peptide, which in turn allows the essential substrate-induced movement of the small domain.

AN 94043333 MEDLINE

DN 94043333 PubMed ID: 8227035

TI Functional roles of **valine** 37 and glycine 38 in the mobile loop of porcine cytosolic **aspartate aminotransferase**.

AU Pan Q W; Tanase S; Fukumoto Y; Nagashima F; Rhee S; Rogers P H; Arnone A; Morino Y

CS Department of Biochemistry, Kumamoto University School of Medicine, Japan.

NC GM-40852 (NIGMS)

SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1993 Nov 25) 268 (33) 24758-65.  
Journal code: HIV; 2985121R. ISSN: 0021-9258.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199312

ED Entered STN: 19940117

Last Updated on STN: 19980206

Entered Medline: 19931220

L6 ANSWER 7 OF 10 CA COPYRIGHT 2001 ACS

AB Aspartate aminotransferase from E. coli, which had been denatured by guanidinium chloride, refolded and reassembled to active dimers in two distinct phases. The unfolded monomer U collapsed within 20 s to an intermediate I\* that was inactive, fluoresced more strongly than, but had the same peptide CD signal as the native dimer. The formation of crosslinkable dimers, as well as the recovery of enzyme activity, occurred

with a biphasic progress curve which was independent of protein concn. The half-lives of the two phases were 100 s and 2000 s. The data are consistent with a three-step mechanism, in which the overall rate of reassembly is detd. by an isomerization of I\* to the assembly-competent monomer M. The latter does not accumulate because it dimerizes rapidly to the active enzyme (D). Reassembly of the enzyme from the compact intermediate M\*, which is **stable** at 1.0M guanidinium chloride, also proceeded in a rapid and a slow phase. Moreover, the formation of

M\*

from the unfolded state was rapid, whereas its refolding to the native dimer was slow. Both the transient intermediate I\* and the equil. intermediate M\* qualify as collapsed intermediate or molten globule states.

AN 116:230721 CA

TI Collapsed intermediates in the reconstitution of dimeric aspartate aminotransferase from Escherichia coli

AU Leistler, Bernd; Herold, Marzell; Kirschner, Kasper

CS Dep. Biophys. Chem., Univ. Basel, Basel, CH-4056, Switz.

SO Eur. J. Biochem. (1992), 205(2), 603-11

CODEN: EJBICA; ISSN: 0014-2956

DT Journal

LA English

L6 ANSWER 8 OF 10 CA COPYRIGHT 2001 ACS

AB The effect of i.m. administration of hydrocortisone (I) [50-23-7] (10 mg/day per animal) for 5 days on the content of the amino acids belonging to the glutamate family in different regions of the mouse brain, along with the activities of glutamine synthetase [9023-70-5], glutamate dehydrogenase [9029-12-3], and **aspartate aminotransferase** [9000-97-9], **alanine aminotransferase** [9000-86-6], tyrosine aminotransferase [9014-55-5] and ornithine [70-26-8] was studied. The activity of **proline** oxidase [9029-17-8] was also studied in these regions. The activities of Na<sup>+</sup>, K<sup>+</sup>-ATPase [9000-83-3] together with the content

of

RNA and protein were also estd. A decrease in the amino acids of the glutamate family in all 3 regions was obsd. with an increase in glutamate dehydrogenase activity in the cerebral cortex. A decrease in the protein content was also obsd., mainly in the brain stem. An increase in Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was obsd. in all 3 regions, with the greatest increase occurring in the cerebral cortex. Apparently, I triggers increased utilization of glutamate in brain as an alternative to glucose, thereby shifting N metab. toward catabolism. The increased activity of Na<sup>+</sup>, K<sup>+</sup>-ATPase under these conditions would further aggravate the same and may lead to membrane **stabilization**.

AN 88:45451 CA

TI Metabolic effects of hydrocortisone in mouse brain

AU Sadasivudu, B.; Rao, T. Indira; Radha, C.; Murthy, Krishna

CS Dep. Biochem., Kurnool Med. Coll., Kurnool, India

SO Neurochem. Res. (1977), 2(5), 521-32

CODEN: NEREDZ

DT Journal

LA English

L6 ANSWER 9 OF 10 MEDLINE

AN 68130568 MEDLINE

DN 68130568 PubMed ID: 5688916

TI Interaction of rat liver **alanine aminotransferase** with **L-proline**.

AU Segal H L; Abraham G J; Matsuzawa T

SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1968 Jan 11) 30 (1) 63-8.  
 Journal code: 9Y8; 0372516. ISSN: 0006-291X.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 196804  
 ED Entered STN: 19900101  
 Last Updated on STN: 19980206  
 Entered Medline: 19680410

L6 ANSWER 10 OF 10 CA COPYRIGHT 2001 ACS  
 AB Expts. were done on **alanine aminotransferase** (I). Rat liver I and I from other organs differed in activity from testis I. Enzyme activity of I was detd. as a function of temp. In the temp. range 55-65.degree., an unexplained break in the curve after .apprx.75% inactivation was regularly observed. First-order decay consts. were obtained in the 1st part of the curve. In protecting liver I against inactivation, arginine, lysine, and ornithine were without effect. Other amino acids were protective, **proline** being the most protective. The prepn. of I from various organs is described. At pH 5.0, 30% (NH4)2SO4 pptd. liver I and also lung, heart, diaphragm, and skeletal muscle I. For pptn. of testis I, 50% concns. of (NH4)2SO4 were required. Antisera were produced in rabbits by injecting liver I. In immunochem. test, the antisera gave a single pptn. band (line of identity) for liver I and for I from diaphragm, skeletal muscle, heart, and lung. Testis I did not react with antisera against liver I. The I from all tissues migrated at the same rate as liver I in acrylamide gel electrophoresis with the exception of testis I, which migrated at .apprx.1/2 the rate. Detns. (in triplicate) of the extinction coeff. of liver I at the absorbancy peak, 278 m.mu., indicated an av. sp. activity of 388 units/mg. protein for liver I and a pyridoxal-P/mole value of 2.0. The results demonstrated that liver I and testis are distinctly different.

I

AN 70:84593 CA  
 TI **Stability** of rat liver alanine aminotransferase and forms of the enzyme in other tissues  
 AU Segal, Harold L.; Abraham, George J.; Schatz, Lillian  
 CS State Univ. of New York, New York, N. Y., USA  
 SO Symp. Pyridoxal Enzymes, 3rd (1968), Meeting Date 1967, 37-42.  
 Editor(s): Yamada, Kozo. Publisher: Maruzen Co. Ltd., Tokyo, Japan.  
 CODEN: 20KFAV  
 DT Conference  
 LA English

Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:sssptal815mxw  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 3 Feb 06 Engineering Information Encompass files have new names  
NEWS 4 Feb 16 TOXLINE no longer being updated  
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure  
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA

NEWS EXPRESS April 18 CURRENT WINDOWS VERSION IS V6.0,  
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),  
AND CURRENT DISCOVER FILE IS DATED 04/06

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 15:02:10 ON 26 APR 2001

=> file ca, biosis, medline  
COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
0.15	0.15

FILE 'CA' ENTERED AT 15:02:35 ON 26 APR 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 15:02:35 ON 26 APR 2001  
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'MEDLINE' ENTERED AT 15:02:35 ON 26 APR 2001

=> s (ASPARTATE AMINOTRANSFERASE?) OR (ALANINE AMINOTRANSFERASE?)  
L1 34077 (ASPARTATE AMINOTRANSFERASE?) OR (ALANINE AMINOTRANSFERASE?)

=> s valine  
L2 65762 VALINE



=> s 11 (p) 12  
L3 139 L1 (P) L2

=> s stab?  
L4 1677944 STAB?

=> s 13 and 14  
L5 6 L3 AND L4

=> dup rem 15  
PROCESSING COMPLETED FOR L5  
L6 4 DUP REM L5 (2 DUPLICATES REMOVED)

=> d 1-4 ab,bib

L6 ANSWER 1 OF 4 CA COPYRIGHT 2001 ACS

AB **Aspartate aminotransferase** or **alanine aminotransferase** in liq. media such as serum or buffered soln. is **stabilized** by adding **valine**, proline, or both into the media to assure the accuracy of clin. studies. The enzymes can be further

**stabilized** by adding a sol. protein such as albumin or gelatin into the media.

AN 131:319672 CA

TI **Stabilization** of aminotransferase using amino acids

IN Baba, Toshiyuki; Tabata, Hiromasa; Nagamatsu, Katashi; Watazu, Yoshifumi; Aoki, Ryoji

PA International Reagents Corporation, Japan; Asahi Chemical Industry Co., Ltd.

SO PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9955850	A1	19991104	WO 1999-JP2205	19990423
	W: CN, JP, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				

PRAI JP 1998-131159 19980424

RE.CNT 9

RE

(2) Beckman Instruments Inc; EP 16573 A CA

(3) Beckman Instruments Inc; US 4325832 A CA

(4) Beckman Instruments Inc; JP 55-141194 A 1980 CA

(5) Iwan Endore Modorobich; US 4652524 A CA

(6) Iwan Endore Modorobich; EP 49475 A CA

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 4 CA COPYRIGHT 2001 ACS

DUPLICATE 1

AB The adiabatic compressibility (.hivin..beta.s) was detd., by the precise sound velocity and d. measurements, for a series of single amino acid substituted mutant enzymes of Escherichia coli dihydrofolate reductase (DHFR) and **aspartate aminotransferase** (AspAT). Interestingly, the .hivin..beta.s values of both DHFR and AspAT were influenced markedly by the mutations at glycine-121 and **valine** -39, resp., in which the magnitude of the change was proportional to the enzyme activity. This result demonstrates that the local change of the primary structure plays an important role in at. packing and protein

dynamics, which leads to the modified **stability** and enzymic function. This is the first report on the compressibility of mutant proteins.

AN 124:224204 CA  
TI A large compressibility change of protein induced by a single amino acid substitution  
AU Gekko, Kunihiro; Tamura, Youjiro; Ohmae, Eiji; Hayashi, Hideyuki; Kagamiyama, Hiroyuki; Ueno, Hiroshi  
CS Fac. Science, Hiroshima Univ., Higashi-Hiroshima, 739, Japan  
SO Protein Sci. (1996), 5(3), 542-5  
CODEN: PRCIEI; ISSN: 0961-8368  
DT Journal  
LA English

L6 ANSWER 3 OF 4 CA COPYRIGHT 2001 ACS

AB A notable feature of porcine cytosolic aspartate aminotransferase is the closure of the active site cleft by a mobile amino-terminal segment (residues 15-40) upon binding substrate. The functional roles of Val17 and Phe18, residues that are part of the mobile loop, have been studied

in the site-directed mutants in which the size and hydrophobic nature of these residues have been changed. Absorption, CD spectra, susceptibility to protease 401, and thermal **stability** did not differ appreciably between wild type and mutant enzymes. In the overall transamination between aspartate and 2-oxoglutarate, V17A represented a typical Km mutant while V17I retained the substrate binding affinity fairly well. In contrast, replacement of Phe18 by Ala resulted in a large

decrease in both catalytic rate and binding affinity for substrates. F18W, F18Y, and F18H showed a moderate decrease in kcat and a

considerable increase in Km values. Single-turnover reactions with four individual substrates yielded analogous results to those obtained for the overall reaction and, in addn., revealed that k/Kd values of mutants F18A and

F18H were over 10 times lower for C5 substrates (glutamate and 2-oxoglutarate) than those for C4 substrates (aspartate and oxalacetate). All mutant enzymes showed variously increased Kd values for substrate analogs such

as 2-methylaspartate, succinate, and glutarate. 1H NMR observations of F18H,

in which His18 served as a built-in probe, were in accord with the behavior that would be expected from the conformational transition. The authors conclude that, although Val17 and Phe18 may not be essential for catalysis, the presence of a bulky residue of appropriate size at each position is crit. for productive binding of substrate.

AN 121:173879 CA  
TI Functional role of the amino-terminal mobile segment in catalysis by porcine cytosolic aspartate aminotransferase. Critical importance of Val17

and Phe18 for productive binding of substrates

AU Nishimura, Kosuke; Higaki, Tsuyoshi; Okamura, Hitoshi; Tanase, Sumio  
CS Sch. Med., Kumamoto Univ., Kumamoto, 860, Japan  
SO J. Biol. Chem. (1994), 269(40), 24712-18  
CODEN: JBCHA3; ISSN: 0021-9258

DT Journal  
LA English

L6 ANSWER 4 OF 4 MEDLINE

AB The functional roles of Val37 and Gly38 in porcine cytosolic aspartate

aminotransferase have been studied in the site-directed mutants V37A, G38A, and G38S where the size and hydrophobic character of these residues has been altered. Previous x-ray studies have shown that Val37 and Gly38, which are part of a flexible loop, interact directly with bound substrate.

From x-ray and solution experiments we find that the V37A, G38A, and G38S mutations do not cause significant perturbations to the unliganded enzyme.

Replacing Val37 with a less bulky alanine residue does not affect the maximal catalytic rate (kcat), but it does increase significantly the Michaelis constants for substrates in the overall transamination reaction between aspartate and 2-oxoglutarate. On the other hand, replacing Gly38 with alanine or serine results in striking decreases in kcat to 5 and 0.6%, respectively, of the value observed for the wild-type enzyme, as well as in considerable increases in Km values. Consequently, the catalytic competence, kcat/Km, decreases by 3 orders of magnitude for

G38A

and by 4 orders of magnitude for G38S. Single turnover reactions of G38A and G38S with four individual substrates (aspartate, glutamate, oxalacetate, and 2-oxoglutarate) are characterized by kinetic parameters that are largely consistent with those of the overall reaction. In addition, the mutations at position 38 impair more seriously the catalytic

competence of the enzyme toward C5-substrates than toward C4-substrates. We conclude that Gly38 is probably required for proper function of the enzyme because it permits a high level of flexibility for the 36-39 peptide, which in turn allows the essential substrate-induced movement of the small domain.

AN 94043333 MEDLINE

DN 94043333 PubMed ID: 8227035

TI Functional roles of **valine** 37 and glycine 38 in the mobile loop of porcine cytosolic **aspartate aminotransferase**.

AU Pan Q W; Tanase S; Fukumoto Y; Nagashima F; Rhee S; Rogers P H; Arnone A; Morino Y

CS Department of Biochemistry, Kumamoto University School of Medicine, Japan.

NC GM-40852 (NIGMS)

SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1993 Nov 25) 268 (33) 24758-65.  
Journal code: HIV; 2985121R. ISSN: 0021-9258.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199312

ED Entered STN: 19940117

Last Updated on STN: 19980206

Entered Medline: 19931220

=>

=>

Connection closed by remote host